



Equivalent boundary conditions for acoustic media with exponential densities

Juliette Chabassier, Marc Duruflé, Victor Péron

► To cite this version:

Juliette Chabassier, Marc Duruflé, Victor Péron. Equivalent boundary conditions for acoustic media with exponential densities. Eighth Singular Days , Jun 2016, Nancy, France. hal-01691768

HAL Id: hal-01691768

<https://inria.hal.science/hal-01691768>

Submitted on 24 Jan 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Equivalent boundary conditions for acoustic media with exponential densities

J. CHABASSIER^{1,2}, M. DURUFLÉ^{3,1}, V. PÉRON^{2,1}

¹ INRIA Bordeaux Sud-Ouest, Team MAGIQUE 3D

² Université de Pau et des Pays de l'Adour

³ ENSEIRB-MATMECA (IPB)

Eighth Singular Days, Nancy

June 26-30, 2016




Introduction

A 3-D Problem set in heterogeneous acoustic media

We solve the Helmholtz equation (with a Dirichlet B.C.) :

$$-\operatorname{div} \left(\frac{1}{\rho} \nabla u \right) - \frac{\omega^2}{\rho c^2} u = f \quad \text{in } \Omega$$



- $\Omega_- = \{r < r_t\}$: Density ρ^- , Velocity c^-
- $\Sigma = \{r = r_t\}$: Interface
- $\Omega_+ = \Omega \setminus \overline{\Omega_-}$: Density ρ^+ , Velocity c^+

Framework

Data and right-hand side

Assumption

- (i) The density ρ and the velocity c are **smooth** and strictly **positive** functions
- (ii) The density ρ^+ satisfies

$$\rho^+(r) = \gamma \exp(-\alpha(r - r_t))$$

where *the parameter α is large*

- (iii) The domain Ω_+ represents a **dissipative medium** :


$$\operatorname{Im} \omega^2 \neq 0 \quad \text{in} \quad \Omega_+$$

Assumption

The right-hand side f has a support in Ω_-

Context


A **boundary layer phenomenon** occurs when α is large : rapid decay of acoustic fields (pressure and velocity) inside the medium Ω_+



Difficulty : Applying a FEM on a mesh with **thin cells** in the **boundary layer** and much larger **elsewhere**

Approach : an Asymptotic Method

- 1 Describing the **boundary layer phenomenon** with a multiscale expansion in power series of α^{-1}
- 2 Deriving Equivalent Boundary Conditions on Σ to replace the **boundary layer** inside Ω_+
- 3 Applying a Finite Element Method to solve the acoustic equation in Ω_- with an Equivalent Boundary Condition



References on **skin effect** in electromagnetism



S. M. RYTOV

Calcul du skin effect par la méthode des perturbations.

Journal of Physics (1940).



E. STEPHAN

Solution procedures for interface problems in [...] electromagnetics.

CISM Courses and Lectures. **277** 291–348 (1983).



R.C MACCAMY, E. STEPHAN

A skin effect approximation for eddy current problems.

Arch. Rational Mech. Anal. **90**(1) (1985) 87–98.



H. HADDAR, P. JOLY, H.-N. NGUYEN

Generalized impedance [...] by strongly absorbing obstacles [...].

Math. Models Methods Appl. Sci. **15**(8) (2005) 1273–1300.



G. CALOZ, M. DAUGE, E. FAOU, V. PÉRON

On the influence of the geometry on skin effect in electromagnetism.

Comput. Methods Appl. Mech. Engrg. 200 (2011), no. 9-12, 1053–1068.

Outline

- 1 **Uniform Estimates**
- 2 **Asymptotic Expansion**
- 3 **Equivalent Conditions**
- 4 **Numerical Results**

Outline


- 1 **Uniform Estimates**
- 2 Asymptotic Expansion
- 3 Equivalent Conditions
- 4 Numerical Results

Problem (P_α)

Helmholtz equation :

$$-\operatorname{div} \left(\frac{1}{\rho} \nabla u_\alpha \right) - \frac{\omega^2}{\rho c^2} u_\alpha = f \quad \text{in } \Omega$$

with an homogeneous Dirichlet boundary condition on $\partial\Omega$.



Here $\rho^+(r) = \gamma \exp(-\alpha(r - r_t))$ and $\alpha > 0$.

Problem (P_α)

Issue : *Uniform H^1 estimates* for solutions u_α of (P_α) w.r.t. $\alpha > 0$?

Problem (\mathbf{P}_α)

Issue : *Uniform H^1 estimates for solutions u_α of (\mathbf{P}_α) w.r.t. $\alpha > 0$?*

Assumption (SA)

The angular frequency ω is not an eigenfrequency of the problem

$$\begin{cases} \operatorname{div} \left(\frac{1}{\rho} \nabla u^- \right) + \frac{\omega^2}{\rho c^2} u^- = 0 & \text{in } \Omega_- \\ u^- = 0 & \text{on } \Sigma \end{cases}$$

Problem (\mathbf{P}_α)

Issue : *Uniform H^1 estimates for solutions u_α of (\mathbf{P}_α) w.r.t. $\alpha > 0$?*

Assumption (SA)

The angular frequency ω is not an eigenfrequency of the problem

$$\begin{cases} \operatorname{div} \left(\frac{1}{\rho} \nabla u^- \right) + \frac{\omega^2}{\rho c^2} u^- = 0 & \text{in } \Omega_- \\ u^- = 0 & \text{on } \Sigma \end{cases}$$

Theorem

Under Assumption (SA), for all $\alpha > 0$ the problem (\mathbf{P}_α) with a right-hand side $f \in L^2(\Omega)$ has a unique solution $u_\alpha \in H_0^1(\Omega)$

$$\|u_\alpha^-\|_{1,\Omega_-} + \|\frac{1}{\sqrt{\rho_+}} u_\alpha^+\|_{0,\Omega_+} + \|\frac{1}{\sqrt{\rho_+}} \nabla u_\alpha^+\|_{0,\Omega_+} \leq C \|f\|_{0,\Omega}$$

Application: Convergence of asymptotic expansion for large parameter α

Problem (P_δ)

Small parameter δ :

$$\delta = \frac{1}{\alpha} \longrightarrow 0 \quad \text{when} \quad \alpha \rightarrow \infty$$

Problem (\mathbf{P}_δ)

Small parameter δ :

$$\delta = \frac{1}{\alpha} \longrightarrow 0 \quad \text{when} \quad \alpha \rightarrow \infty$$

Problem (\mathbf{P}_δ) writes :

$$-\operatorname{div}\left(\frac{1}{\rho}\nabla u_\delta^-\right) - \frac{\omega^2}{\rho c^2}u_\delta^- = f \quad \text{in } \Omega_-$$

$$-\Delta u_\delta^+ - \frac{1}{\delta}\partial_r u_\delta^+ - \frac{\omega_+^2}{c^2}u_\delta^+ = 0 \quad \text{in } \Omega_+$$

$$u_\delta^+ = u_\delta^- \quad \text{on } \Sigma$$

$$\partial_{\mathbf{n}} u_\delta^+ = \partial_{\mathbf{n}} u_\delta^- \quad \text{on } \Sigma$$

$$u_\delta^+ = 0 \quad \text{on } \partial\Omega$$

Outline

- 1 Uniform Estimates
- 2 Asymptotic Expansion**
- 3 Equivalent Conditions
- 4 Numerical Results

Asymptotic Expansion

Overview

Deriving an Asymptotic Expansion for the solution (u_δ^-, u_δ^+) of (\mathbf{P}_δ) when $\delta \rightarrow 0$:

$$u_\delta^-(\mathbf{x}) = u_0^-(\mathbf{x}) + \delta u_1^-(\mathbf{x}) + \delta^2 u_2^-(\mathbf{x}) + \delta^3 u_3^-(\mathbf{x}) + \cdots \quad \text{in } \Omega_-$$

$$u_\delta^+(\mathbf{x}) = u_0^+(\mathbf{x}; \delta) + \delta u_1^+(\mathbf{x}; \delta) + \delta^2 u_2^+(\mathbf{x}; \delta) + \delta^3 u_3^+(\mathbf{x}; \delta) + \cdots \quad \text{in } \Omega_+,$$

$$\text{with } u_j^+(\mathbf{x}; \delta) = \chi(r) \mathfrak{U}_j \left(\theta, \phi, \frac{r - r_t}{\delta} \right)$$

Method based on the Scaling :

$$S = \frac{r - r_t}{\delta}$$

Equations for the terms \mathfrak{U}_n and u_n^-

$$\begin{cases} -\partial_S^2 \mathfrak{U}_n - \partial_S \mathfrak{U}_n &= \sum_{p=1}^n A_p \mathfrak{U}_{n-p} & \text{in } \Sigma \times (0, +\infty), \\ \partial_S \mathfrak{U}_n &= \partial_n u_{n-1}^- & \text{on } \Sigma, \end{cases}$$

and

$$\begin{cases} -\operatorname{div}\left(\frac{1}{\rho} \nabla u_n^-\right) - \frac{\omega^2}{\rho c^2} u_n^- &= f \delta_n^0 & \text{in } \Omega_-, \\ u_n^- &= \mathfrak{U}_n & \text{on } \Sigma. \end{cases}$$

First terms

1

$$\mathfrak{U}_0 = 0$$

2

$$\begin{cases} -\operatorname{div}\left(\frac{1}{\rho}\nabla u_0^-\right) - \frac{\omega^2}{\rho c^2}u_0^- = f & \text{in } \Omega_- \\ u_0^- = 0 & \text{on } \Sigma \end{cases}$$

3

$$\mathfrak{U}_1(\cdot, s) = -\partial_{\mathbf{n}}u_0^- e^{-s}, \quad s \in (0, \infty)$$

4

$$\begin{cases} -\operatorname{div}\left(\frac{1}{\rho}\nabla u_1^-\right) - \frac{\omega^2}{\rho c^2}u_1^- = 0 & \text{in } \Omega_- \\ u_1^- = -\partial_{\mathbf{n}}u_0^- & \text{on } \Sigma \end{cases}$$

Higher order terms

•

$$\mathfrak{U}_2(\cdot, S) = (a_2 + b_2 S) e^{-S}, \quad S \in (0, \infty)$$

Here

$$\begin{cases} a_2 = -\partial_{\mathbf{n}} u_1^- + \frac{2}{r_t} \partial_{\mathbf{n}} u_0^- , \\ b_2 = \frac{2}{r_t} \partial_{\mathbf{n}} u_0^- . \end{cases}$$

•

$$\begin{cases} -\operatorname{div}\left(\frac{1}{\rho} \nabla u_2^-\right) - \frac{\omega^2}{\rho c^2} u_2^- = 0 & \text{in } \Omega_- , \\ u_2^- = -\partial_{\mathbf{n}} u_1^- + \frac{2}{r_t} \partial_{\mathbf{n}} u_0^- & \text{on } \Sigma . \end{cases}$$

Higher order terms



$$\mathfrak{L}_3(\theta, \phi, S) = (a_3(\theta, \phi) + Sb_3(\theta, \phi) + S^2c_3(\theta, \phi))e^{-S}, \quad S \in (0, \infty)$$

Here

$$\begin{cases} a_3 = -\partial_{\mathbf{n}}u_2^- + \frac{2}{r_t}\partial_{\mathbf{n}}u_1^- - \left\{ \frac{6}{r_t^2} + \frac{\omega_+^2}{c^2} + \Delta_\Sigma \right\} \partial_{\mathbf{n}}u_0^-, \\ b_3 = \frac{2}{r_t}\partial_{\mathbf{n}}u_1^- - \left\{ \frac{6}{r_t^2} + \frac{\omega_+^2}{c^2} + \Delta_\Sigma \right\} \partial_{\mathbf{n}}u_0^-, \\ c_3 = -\frac{3}{r_t^2}\partial_{\mathbf{n}}u_0^-. \end{cases}$$



$$\begin{cases} -\operatorname{div}\left(\frac{1}{\rho}\nabla u_3^-\right) - \frac{\omega^2}{\rho c^2}u_3^- = 0 & \text{in } \Omega_-, \\ u_3^- = -\partial_{\mathbf{n}}u_2^- + \frac{2}{r_t}\partial_{\mathbf{n}}u_1^- - \left\{ \frac{6}{r_t^2} + \frac{\omega_+^2}{c^2} + \Delta_\Sigma \right\} \partial_{\mathbf{n}}u_0^- & \text{on } \Sigma. \end{cases}$$

Existence and regularity of the asymptotics

Proposition

Let $k \in \mathbb{N}$. Assume that $f \in H^k(\Omega_-)$ and $\rho \in C^\infty(\overline{\Omega_-})$. Then it is possible to derive the first $(k + 2)$ terms (u_j^-, \mathfrak{U}_j) , and

$$u_0^- \in H^{k+2}(\Omega_-), u_1^- \in H^{k+1}(\Omega_-), \dots, u_{k+1}^- \in H^1(\Omega_-)$$

and

$$\mathfrak{U}_1 \in H^{k+\frac{1}{2}}(\Sigma \times \mathbb{R}^+), \mathfrak{U}_2 \in H^{k-\frac{1}{2}}(\Sigma \times \mathbb{R}^+), \dots, \mathfrak{U}_{k+1} \in H^{\frac{1}{2}}(\Sigma \times \mathbb{R}^+)$$

Validation of the Asymptotic Expansion

Aim : proving Estimates for Remainders

$$r_\delta^N := u_\delta - \sum_{n=0}^N \delta^n u_n \quad \text{in } \Omega$$

Evaluation of the right hand sides in

$$\left\{ \begin{array}{ll} -\operatorname{div} \left(\frac{1}{\rho} \nabla r_\delta^{N,-} \right) - \frac{\omega^2}{\rho c^2} r_\delta^{N,-} = 0 & \text{in } \Omega_- \\ -\operatorname{div} \left(\frac{1}{\rho} \nabla r_\delta^{N,+} \right) - \frac{\omega_+^2}{\rho c^2} r_\delta^{N,+} = f_{N,\delta} & \text{in } \Omega_+ \\ r_\delta^{N,+} = r_\delta^{N,-} & \text{on } \Sigma \\ \partial_{\mathbf{n}} r_\delta^{N,+} = \partial_{\mathbf{n}} r_\delta^{N,-} + g_{N,\delta} & \text{on } \Sigma \\ r_\delta^N = 0 & \text{on } \partial\Omega \end{array} \right.$$

The RHS are explicit : $f_{N,\delta} = \mathcal{O}(\delta^{N-1})$ and $g_{N,\delta} = \delta^N \partial_{\mathbf{n}} u_N^-$

$$\|r_\delta^{N,-}\|_{1,\Omega_-} + \delta^{-1/2} \left\| \frac{1}{\sqrt{\rho_+}} r_\delta^{N,+} \right\|_{0,\Omega_+} + \delta^{1/2} \left\| \frac{1}{\sqrt{\rho_+}} \nabla r_\delta^{N,+} \right\|_{0,\Omega_+} \leq C_N \delta^{N+1}$$

Outline

- 1 Uniform Estimates
- 2 Asymptotic Expansion
- 3 Equivalent Conditions**
- 4 Numerical Results

Equivalent Conditions on Σ

Overview

We identify a simpler problem satisfied by

$$u_{k,\delta}^- := u_0^- + \delta u_1^- + \delta^2 u_2^- + \cdots + \delta^k u_k^- \quad \text{up to } \mathcal{O}(\delta^{k+1})$$

The simpler problem *of order $k + 1$* writes

$$(\mathbf{P}_\delta^k) \quad \begin{cases} -\operatorname{div}\left(\frac{1}{\rho}\nabla u_k^\delta\right) - \frac{\omega^2}{\rho c^2}u_k^\delta & = f & \text{in } \Omega_-, \\ u_k^\delta + D_{k,\delta}(\partial_{\mathbf{n}}u_k^\delta) & = 0 & \text{on } \Sigma, \end{cases}$$

with $D_{k,\delta}$ a surface differential operator

Equivalent Conditions

- ❶ Order 1 :

$$u_0 = 0 \quad \text{on} \quad \Sigma$$

- ❷ Order 2 :

$$u_1^\delta + \delta \partial_{\mathbf{n}} u_1^\delta = 0 \quad \text{on} \quad \Sigma$$

- ❸ Order 3 :

$$u_2^\delta + \delta \left(1 - \frac{2\delta}{r_t} \right) \partial_{\mathbf{n}} u_2^\delta = 0 \quad \text{on} \quad \Sigma$$

- ❹ Order 4 :

$$u_3^\delta + \delta \left(1 - \frac{2\delta}{r_t} + \delta^2 \left\{ \frac{6}{r_t^2} - \frac{\omega_+^2}{c^2} - \Delta_\Sigma \right\} \right) \partial_{\mathbf{n}} u_3^\delta = 0 \quad \text{on} \quad \Sigma$$

Dirichlet-to-Neumann Conditions

- Order 4 :

$$\delta^{-1} \left(\left(1 + \frac{2\delta}{r_t} \right) \mathbb{I} - \delta^2 \left\{ \frac{2}{r_t^2} + \frac{\omega_+^2}{c^2} + \Delta_\Sigma \right\} \right) v_3^\delta + \partial_{\mathbf{n}} v_3^\delta = 0 \quad \text{on } \Sigma$$

Dirichlet-to-Neumann Conditions

- Order 4 :

$$\delta^{-1} \left(\left(1 + \frac{2\delta}{r_t} \right) \mathbb{I} - \delta^2 \left\{ \frac{2}{r_t^2} + \frac{\omega_+^2}{c^2} + \Delta_\Sigma \right\} \right) v_3^\delta + \partial_{\mathbf{n}} v_3^\delta = 0 \quad \text{on } \Sigma$$

For all $k \in \{1, 2, 3\}$, the simpler problem writes

$$(\mathbf{q}_\delta^k) \quad \begin{cases} -\operatorname{div}\left(\frac{1}{\rho} \nabla v_k^\delta\right) - \frac{\omega^2}{\rho c^2} v_k^\delta &= f \quad \text{in } \Omega_- , \\ N_{k,\delta}(v_k^\delta) + \partial_{\mathbf{n}} v_k^\delta &= 0 \quad \text{on } \Sigma , \end{cases}$$

with $N_{k,\delta}$ a surface differential operator.

Stability and convergence results

Theorem

Under Assumption (SA), there exists $\delta_0 > 0$ s.t. for all $\delta \in (0, \delta_0)$, the problem (\mathbf{Q}_δ^k) with data $f \in L^2(\Omega_-)$ has a unique solution $v_k^\delta \in V_k$, and

$$\|v_k^\delta\|_{1,\Omega_-} \leq C\|f\|_{0,\Omega_-}$$

$$\|u_\delta - v_k^\delta\|_{1,\Omega_-} \leq C_k \delta^{k+1}$$

Functional setting : $v_k^\delta \in V_k$ where

$$V_k = H^1(\Omega_-) \quad \text{when } k = 1, 2$$

and

$$V_3 = \{u \in H^1(\Omega_-) \mid u|_\Sigma \in H^1(\Sigma)\}$$

Proof of the stability result

Lemma

Under Assumption (SA), there exists $\delta_0 > 0$ s.t. for all $\delta \in (0, \delta_0)$, any solution $v_k^\delta \in V_k$ of problem (\mathbf{Q}_δ^k) with a data $f \in L^2(\Omega_-)$ satisfies

$$\|v_k^\delta\|_{0,\Omega_-} \leq C \|f\|_{0,\Omega_-} .$$

Sketch of the proof of the Lemma

Assume : there exists $(u_m) \in (V_k)^{\mathbb{N}}$ satisfying $(Q_{\delta_m}^k)$ with $\delta_m \rightarrow 0$ and $f_m \in L^2(\Omega_-)$ such that

$$\|u_m\|_{0,\Omega_-} = 1 \quad \text{and} \quad \|f_m\|_{0,\Omega_-} \rightarrow 0$$

We prove successively that :

1

$$\|u_m\|_{1,\Omega_-} \leq C$$

2 there exists a subsequence (u_m) s.t.

$$u_m \rightarrow u \quad \text{in} \quad L^2(\Omega_-) \quad \text{and} \quad \nabla u_m \rightharpoonup \nabla u \quad \text{in} \quad L^2(\Omega_-)$$

$$\text{and } u_m \rightarrow 0 \quad \text{in} \quad L^2(\Sigma)$$

3

$$\|u\|_{0,\Omega_-} = 1$$

4 Using Assumption (SA), we prove that $u = 0$. Contradiction.

Outline

- 1 Uniform Estimates
- 2 Asymptotic Expansion
- 3 Equivalent Conditions
- 4 Numerical Results**

Numerical method

Helmholtz equation with radial data

Using a decomposition in spherical harmonics we solve a sequence of 1-D problems with a Finite Element Method.

- 1-D Computational domain :

Approximate models : $\Omega_- = (0; 1)$

Reference solutions : $\Omega = (0; R_\Omega)$ such that $\rho^+(R_\Omega) = 10^{-15}$

- \mathbb{P}_{10} -finite elements (Lagrange) available in the Library Montjoie
- Data :
 - 1 Artificial data
 - 2 Realistic data

Artificial data

Framework

- The density is given as

$$\rho(r) = \begin{cases} D_1 & \text{if } r < 1 \\ D_1 e^{-(r-1)/\delta} & \text{otherwise} \end{cases}$$

$$D_1 = 2 \times 10^{-4} \text{ kg/m}^3,$$

- The velocity is constant


$$c = 8 \times 10^3 \text{ m/s}$$

- **Radial gaussian source** located at the radius $r = 0.8$
- $\omega = 2\pi f_0$ where $f_0 \in \{7 \times 10^4 \text{ Hz}, 7 \times 10^5 \text{ Hz}\}$

Convergence results

Freq. $f_0 = 7 \times 10^4$ Hz


Relative L^2 -error between exact solution and approximate solutions versus δ



Convergence results

Freq. $f_0 = 7 \times 10^5$ Hz

Relative L^2 -error between exact solution and approximate solution versus δ



Realistic data

Using the data given by the Standard Solar Model



CHRISTENSEN-DALSGAARD ET AL, THE CURRENT STATE OF SOLAR MODELING. SCIENCE, 1996





Figure : Exact value and approximated value of ρ^{sun} versus the relative radius r/R_{\odot} .

Realistic data

Reference solution in the plane Oxz

Real part of $u/\sqrt{\rho}$ when $\delta = 1/7000$




The source is a 3-D gaussian and we use 100 spherical harmonics

Realistic setting


Convergence results

Relative L^2 -error between reference solution and approximate solution versus δ



Realistic setting


Comparison of equivalent boundary conditions for different frequencies from 0.3 mHz until 7 mHz.



Thank you for your attention

Numerical tests with artificial data

Qualitative Comparisons




Reference solution and solutions using equivalent boundary conditions.

$$f_0 = 7 \times 10^4 \text{ Hz}$$

Numerical tests with artificial data

Qualitative Comparisons



Reference solution and solutions using atmosphere boundary conditions.

$$f_0 = 7 \times 10^5 \text{ Hz}$$

Realistic setting

- The density is given as

$$\rho_0(r) = \left(1 + \frac{2i\gamma}{\omega}\right) \begin{cases} \rho^{\text{sun}} & \text{if } r/R_{\odot} \leq 1.0007126 \\ D_1 e^{-(r/R_{\odot}-1)/\delta} & \text{otherwise} \end{cases}$$

$$D_1 \approx 3.292 \times 10^{-6} \text{ kg/m}^3, \gamma = \frac{\omega}{100}$$

- The velocity is given as

$$c_0(r) = \begin{cases} c^{\text{sun}} & \text{if } r/R_{\odot} \leq 1.0007126 \\ C_1 & \text{otherwise} \end{cases}$$

$$C_1 \approx 6.865 \times 10^3 \text{ m/s}$$

- The source is a 3-D gaussian:

$$f(x) = \exp(-\beta ||x - x_0||^2)$$

Here $x_0 \approx (0, 0, 0.981)$ and β is chosen such that $\exp(-0.015 \beta) = 10^{-6}$.